



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number:

0 665 013 A1

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: **95100194.0**

(51) Int. Cl.⁶: **A61K 31/42, A61K 31/275**

(22) Date of filing: **09.01.95**

(30) Priority: **13.01.94 US 181116**

(43) Date of publication of application:
02.08.95 Bulletin 95/31



(84) Designated Contracting States:
**AT BE CH DE DK ES FR GB GR IE IT LI LU NL
PT SE**

(71) Applicant: **HOECHST AKTIENGESELLSCHAFT**
Brüningstrasse 50
D-65929 Frankfurt am Main (DE)

(72) Inventor: **Bartlett, Robert Ryder, Dr.**
Schmittweg 23
D-64291 Darmstadt (DE)
Inventor: **Morris, Randall E.**
1765 Fallen Leaf Lane
Los Altos, CA 94024 (US)

(54) **Method of treating hyperproliferative vascular disease.**

(57) A method of preventing or treating hyperproliferative vascular disease in a mammal consists of administering an effective amount of a compound of the formula I or II



to said mammal.

EP 0 665 013 A1

Best Available Copy

European Patent 13,376 discloses N-(4-trifluoromethylphenyl)-5-methylisoxazole-4-carboxamide (compound 1) as being anti-inflammatory. Processes for the preparation of this compound are also described therein.

It is additionally known that the compound 1 and N-(4-trifluoromethylphenyl)-2-cyano-3-hydroxy-ycrotonamide (compound 2) have immunomodulation properties, so that they are suitable as pharmaceutical against chronic graft versus host diseases and against autoimmune disorders, in particular systemic Lupus erythematosus (EP 0,217,206).

US Patent 4,061, 767 describes the use of 2-hydroxyethylidene-cyanoacetanilide derivatives for the preparation of pharmaceuticals having anti-inflammatory and analgesic action.

Many individuals suffer from heart disease caused by a partial blockage of the blood vessels that supply the heart with nutrients. More severe blockage of blood vessels in such individuals often leads to hypertension, ischemic injury, stroke, or myocardial infarction. Typically vascular occlusion is preceded by vascular stenosis resulting from intimal smooth muscle cell hyperplasia. The underlying cause of the intimal smooth muscle cell hyperplasia is vascular smooth muscle injury and disruption of the integrity of the endothelial lining. The overall disease process can be termed a hyperproliferative vascular disease because of the etiology of the disease process. Intimal thickening following arterial injury can be divided into three sequential steps: 1) initiation of smooth muscle cell proliferation following vascular injury, 2) smooth muscle cell migration to the intima, and 3) further proliferation of smooth muscle cells in the intima with deposition of matrix. Investigations of the pathogenesis of intimal thickening have shown that, following arterial injury, platelets, endothelial cells, macrophages and smooth muscle cells release paracrine and autocrine growth factors (such as platelet derived growth factor, epidermal growth factor, insulin-like growth factor, and transforming growth factor) and cytokines that result in the smooth muscle cell proliferation and migration. T-cells and macrophages also migrate into the neointima. This cascade of events is not limited to arterial injury, but also occurs following injury to veins and arterioles.

Vascular injury causing intimal thickening can be broadly categorized as being either biologically or mechanically induced. Atherosclerosis is one of the most commonly occurring forms of biologically mediated vascular injury leading to stenosis. The migration and proliferation of vascular smooth muscle plays a crucial role in the pathogenesis of atherosclerosis. Atherosclerotic lesions include massive accumulation of lipid laden "foam cells" derived from monocyte/macrophage and smooth muscle cells. Formation of "foam cell" regions is associated with a breach of endothelial integrity and basal lamina destruction. Triggered by these events, restenosis is produced by a rapid and selective proliferation of vascular smooth muscle cells with increased new basal lamina (extracellular matrix) formation and results in eventual blocking of arterial pathways.



Mechanical injuries leading to intimal thickening result following balloon angioplasty, vascular surgery, transplantation surgery, and other similar invasive processes that disrupt vascular integrity. Intimal thickening following balloon catheter injury has been studied in animals as a model for arterial restenosis that occurs in human patients following balloon angioplasty. It has been shown that deendothelization with an intraarterial catheter that dilates an artery injures the innermost layers of medial smooth muscle and may even kill some of the innermost cells. Injury is followed by a proliferation of the medial smooth muscle cells, after which many of them migrate into the intima through fenestrae in the internal elastic lamina and proliferate to form a neointimal lesion.

Vascular stenosis can be detected and evaluated using angiographic or sonographic imaging techniques and is often treated by percutaneous transluminal coronary angioplasty (balloon catheterization). Within a few months following angioplasty, however, the blood flow is reduced in approximately 30-40 percent of these patients as a result of restenosis caused by a response to mechanical vascular injury suffered during the angioplasty procedure, as described above.

In an attempt to develop better agents for preventing or reducing smooth muscle proliferation and intimal thickening, the use of the balloon catheter induced arterial injury in a variety of mammals has been developed as a standard model of vascular injury that will lead to intimal thickening and eventual vascular narrowing.

Surprisingly a compound of the formula I or II shows an effective inhibition of intimal thickening in injured carotid arteries.

Therefore the invention relates to a method of preventing or treating hyperproliferative vascular disease in a mammal in need thereof by administering an amount, effective to inhibit intimal thickening in said mammal, of a compound of the formula I or II



wherein R¹ denotes

- a) methyl,
- b) (C₃-C₆)-cycloalkyl,
- c) (C₂-C₆)-alkyl, having 1 or 2 triple bonds between the carbon atoms,

R² denotes

- a) -CF₃ or
- b) -CN,

R³ denotes

- a) (C₁-C₄)-alkyl or
- b) hydrogen atom,

X denotes

- a) carbon atom or
- b) nitrogen atom,

the compound of the formula II being present as such or in the form of a physiologically tolerable salt.

Preferred are compounds of the formula I or II

wherein R¹ denotes

- a) methyl,
- b) cyclopropyl or
- c) -CH₂-CH₂-C≡CH,

R² denotes -CF₃,

R³ denotes methyl or hydrogen atom and


X denotes carbon atom.

Especially preferred is a compound of the formula I or II, wherein R¹ denotes methyl, R² denotes -CF₃, R³ denotes hydrogen atom and X denotes carbon atom.


Suitable physiologically tolerable salts of the compound of the formula II are, for example, alkali metal, alkaline earth metal or ammonium salts, including those of physiologically tolerable organic ammonium bases.

The compounds of the formula I or II can be prepared by the following process:

A compound of the formula III



in which Y represents a halogen atom, preferably chlorine or bromine, is reacted with the amine of the formula (IV)



wherein R^2 , R^3 and X have the same meanings as in formula I, to give the compound of the formula I, and this can then be reacted in the presence of a basic agent to give the compound of the formula II.

The starting substances for the reactions are known or can be easily prepared by methods known from the literature. The above-mentioned reactions are carried out under standard conditions in a known manner (EP 13,376; EP 484,223; EP 538,783; US 4,061,767).

The invention also relates to pharmaceuticals which contain an effective amount of compound of the formula I and/or compound of the formula II, the compound of the formula II being present as such or in the form of a physiologically tolerable salt, in addition to pharmaceutically suitable and physiologically tolerable excipients, diluents and/or other active substances and auxiliaries.

The invention also relates to a process for the preparation of a pharmaceutical for the treatment of hyperproliferative vascular disease, which comprises bringing a compound of the formula I or II or a physiologically tolerable salt of compound of the formula II or a mixture thereof into a suitable administration form using a pharmaceutically suitable and physiologically acceptable excipient and, if appropriate, other suitable active substances, additives or auxiliaries.

The pharmaceutical according to the invention can be administered orally, intravascularly, topically, rectally, parenterally, intranasally, intrabronchially, transdermally or via a vascular stent impregnated with a compound of the formula I or II. Administration is carried out before, during and/or after a biologically or mechanically mediated vascular injury, or under conditions that would predispose a mammal to suffering such a vascular injury.

Suitable solid or liquid pharmaceutical administration forms are, for example, granules, powders, coated tablets, tablets, (micro)capsules, suppositories, syrups, juices, suspensions, emulsions, drops or injectable solutions and preparations having a protracted release of active substance, within these preparations customary auxiliaries, such as excipients, disintegrants, binders, coating agents, swelling agents, lubricants, flavorings, sweeteners or solubilizers are used. Commonly used auxiliaries are, for example, magnesium carbonate, titanium dioxide, lactose, mannitol and other sugars, talc, milk protein, gelatin, starch, cellulose and its derivatives, animal and vegetable oils, polyethylene glycols and solvents, such as, for example, sterile water and mono- or polyhydric alcohols, for example glycerol.

Preferably, the pharmaceutical preparations are prepared and administered in dosage units, each unit containing as the active constituent a certain dose of compound of the formula I and/or II, where compound of the formula II is present as such or in the form of a physiologically tolerable salt. In the case of solid dosage units, such as tablets, capsules or suppositories, this dose can be up to about 500 mg, preferably 5 to 400 mg, more preferably 5 to 200 mg, in particular 10 to 100 mg, and especially 10 to 25 mg.

For the treatment of a patient (70 kg) suffering from a hyperproliferative vascular disease in the early phases an intravenous infusion treatment of at most 1200 mg per day and in the later rehabilitation phases an oral administration of 3 times 300 mg per day of compound of the formula I and/or II and/or of the corresponding salts of compound of the formula II are indicated.

Under certain circumstances, however, higher or lower doses may also be appropriate. The administration of the dose can be carried out both by singly administration in the form of an individual dosage unit or else several smaller dosage units and by multiple administration of subdivided doses at specific intervals.

A compound of the formula I or II and/or its corresponding salts can also be combined during the preparation of the above-mentioned pharmaceutical administration forms together with other immunosuppressive substances, for example rapamycin (Wyeth-Ayerst Inc., Princeton, NJ), mycophenolic acid (Sigma Chemical Co., St. Louis, MO), cyclosporin A (Sandoz Inc., East Hannover, NJ), FK506 (Fujisawa Inc., Osaka, Japan) or brequinar sodium.

As such, the compound of the formula I or II is useful, alone or in combination with other pharmacological effective substances, in preventing or treating intimal smooth muscle cell hyperplasia, restenosis, and vascular occlusion in a mammal, particularly following either biologically or mechanically mediated vascular injury, or under conditions that would predispose a mammal to suffering such a vascular injury. Biologically mediated vascular injury includes, but is not limited to injury to autoimmune disorders; alloimmune related disorders; infectious disorders including endotoxins and herpes viruses such as cytomegalovirus; metabolic disorders such as atherosclerosis; and vascular injury resulting from hypothermia and irradiation. Mechanically

cally mediated vascular injury includes, but is not limited to vascular injury caused by catheterization procedures or vascular scraping procedures such as percutaneous transluminal coronary angioplasty; vascular surgery; transplantation surgery; laser treatment, and other invasive procedures which disrupt the integrity of the vascular intima or endothelium.

- 5 Preventing includes the prophylactic prevention of hyperproliferative vascular disease in a susceptible mammal and treating includes arresting the development, and retarding the progression of hyperproliferative vascular disease in a susceptible mammal.

Example 1

10

Pharmacological tests and results

The effect of N-(4-trifluoromethylphenyl)-5-methylisoxazole-carboxamide (compound 1) on hyperproliferative vascular disease was established in an in vivo standard pharmacological test procedure that emulates the hyperproliferative effects observed in mammals that are undergoing intimal smooth muscle proliferation and are therefore developing restenosis. Cyclosporin A was also evaluated in these test procedure for the purpose of comparison.

Compound 1 was also evaluated in an in vivo standard pharmacological test procedure that emulates the vascular injury suffered and restenosis that develops following percutaneous transluminal coronary angioplasty in humans.

Intimal smooth muscle proliferation was produced by balloon catheter injury to the left carotid artery of groups of 6, male Sprague-Dawley rats. Endothelial denudation and vascular injury were achieved in the left carotid arteries of male Sprague-Dawley rats. A balloon catheter (2 French Fogarty, Edwards Laboratories, Santa Anna, CA) was passed through the external carotid artery into the aorta. The balloon was then inflated with an amount of water sufficient to distend the common carotid artery and was then pulled back to the external carotid artery. The inflation and pull back were repeated three times. This procedure leads to complete denudation of the endothelium throughout the common carotid artery, and also some injury typically occurs to the media smooth muscle cells.

During a 17-days period (3 days before operation to 13 days after operation), these rats were divided into 2 groups of 6 animals and treated daily with compound 1 (20 mg/kg/day, intraperitoneal (i.p.)) or cyclosporin A (3 mg/kg/day; i.p.).

An injured group not treated with any drug was used as an injured control to establish the amount of intimal thickening in the absence of treatment. The right carotid was used as an uninjured control in all groups.

After the 14 day post-operative period, the rats were sacrificed, the carotids removed. The mean areas of the intima, media and total blood vessel wall were measured by morphometry. The injured artery was also examined using histopathologic assays. Results are presented as an intima percent, expressed as follows:

40

area of intima

* 100

area of intima + area of media

45

The following table shows the data obtained in the above experiment.
Effect of test substances on intimal thickening in injured carotid arteries

50

Test Group	Intima Percent (\pm S.E.) ^{*)}
Uninjured Control	0.00 \pm 0.00
Untreated Injured Control	46.06 \pm 16.36
Compound 1 (20 mg/kg)	7.88 \pm 7.88
CsA ^{*)} (3 mg/kg - 14 days)	39.39 \pm 26.06

55

^{*)} S.E. = standard error, CsA = cyclosporin A.


These results show that treatment with compound 1 resulted in an 83 % decrease in the mean percentage intimal thickening compared with the untreated injured control group. Cyclosporin A failed to produce any meaningful reduction in intimal thickening.

The results of the in vivo standard test procedure demonstrate that compound 1 is useful in preventing or treating hyperproliferative vascular disease. Specifically, compound 1 is useful in preventing or treating intimal smooth muscle cell hyperplasia, restenosis, and vascular occlusion in a mammal, particularly following either biologically or mechanically mediated vascular injury, or under conditions that would predispose a mammal to suffering such a vascular injury.

10 Example 2

Preparation of N-(4-trifluoromethylphenyl)-5-methylisoxazole-4-carboxamide

(Compound 1)




A solution of 0.05 mol of 5-methylisoxazole-4-carbonyl chloride (7.3 g) in 20 ml of acetonitrile is added drop wise at room temperature to a solution of 0.1 mol of 4-trifluoromethylaniline (16.1 g) in 150 ml of acetonitrile. After stirring for 20 minutes, the precipitated 4-trifluoromethylaniline hydrochloride is filtered off with suction and washed twice with 20 ml of acetonitrile each time, and the combined filtrates are concentrated under reduced pressure. The yield is 12.8 g of white, crystalline N-(4-trifluoromethylphenyl)-5-methylisoxazole-4-carboxamide (compound 1).

Melting point from toluol 166,5 °C

35 Example 3

Preparation of N-(4-trifluoromethylphenyl)-2-cyano-3-hydroxycrotonamide

(compound 2)



0.1 mol of N-(4-trifluoromethylphenyl)-5-methylisoxazole-4-carboxamide is dissolved in 100 ml of methanol and treated at + 10 °C with a solution of 0.11 mol (4.4 g) of sodium hydroxide in 100 ml of water. The mixture is stirred for 30 minutes and, after diluting with water, is acidified with concentrated hydrochloric acid. The precipitated crop of crystals is filtered off with suction, washed with water and dried in air.

The yield is 24.4 g of N-(4-trifluoromethylphenyl)-2-cyano-3-hydroxycrotonamide (compound 2).

Melting point from methanol 205 to 206 °C.

Example 4

Acute toxicity after intraperitoneal administration



- 5 The acute toxicity after intraperitoneal administration of the test substances was determined with NMRI mice (20 to 25 g) and SD rats (120 to 195 g). The test substance was suspended in a 1 % strength sodium carboxymethylcellulose solution. The different dosages of the test substance were administered to the mice in a volume of 10 ml/kg of body weight and to the rats in a volume of 5 ml/kg of body weight. Per dosage, 10 animals were used. After 3 weeks, the acute toxicity was determined by the method of Litchfield and
 10 Wilcox. The results are summarized in the table 2.

Table 2

15	Compound 1 acute toxicity intraperitoneal LD ₅₀ (mg/kg)	Compound 2 acute toxicity intraperitoneal LD ₅₀ (mg/kg)
NMRI mouse	185 (163 - 210)	150 (100 - 200)
SD rat	170 (153 - 189)	

Claims

1. A method of preventing or treating hyperproliferative vascular disease in a mammal, comprising administering an amount, effective to inhibit intimal thickening in said mammal, of a compound of the
 25 formula I or II



wherein R¹ denotes

- a) methyl,
 b) (C₃-C₆)-cycloalkyl,
 c) (C₂-C₆)-alkyl, having 1 or 2 triple bonds between the carbon atoms,

R² denotes

- a) -CF₃ or
 b) -CN,

R³ denotes

- a) (C₁-C₄)-alkyl or
 b) hydrogen atom,

X denotes

- a) carbon atom or
 b) nitrogen atom,

the compound of formula II being present as such or in the form of a physiologically tolerable salt.

2. The method of claim 1, wherein the compound of the formula I or II is selected from the group comprising where R¹ denotes

- a) methyl,
- b) cyclopropyl or
- c) $-\text{CH}_2-\text{CH}_2-\text{C}\equiv\text{CH}$,

R^2 denotes $-\text{CF}_3$,

R^3 denotes methyl or hydrogen atom and

X denotes carbon atom.

3. The method of claim 1, wherein the compound of the formula I or II is selected from the group comprising N-(4-trifluoromethylphenyl)-2-cyano-3-hydroxycrotonamide and N-(4-trifluoromethylphenyl)-5-methylisoxazole-4-carboxamide.
4. The method of claim 1, wherein the administering is accomplished orally, parenterally, intravascularly, intranasally, intrabronchially, transdermally, rectally, or via a vascular stent impregnated with a compound of the formula I or II.
5. The method of claim 1, wherein the compound of the formula I or II is administered concurrent with said mammal undergoing a percutaneous transluminal coronary angioplasty procedure.
6. The method of claim 5, which further comprises administering the compound of the formula I or II subsequent to said mammal undergoing a percutaneous transluminal coronary angioplasty procedure.
7. The method of claim 1, wherein the hyperproliferative vascular disease is selected from the group consisting of intimal smooth muscle cell hyperplasia, restenosis, and vascular occlusion.
8. The method of claim 1, wherein the compound of the formula I or II is administered prior to, concurrent with or subsequent to said mammal sustaining a biologically mediated vascular injury.
9. The method of claim 1, wherein a compound of the formula I or II is administered prior to, concurrent with or subsequent to said mammal sustaining a mechanically mediated vascular injury.
10. A composition for the use in preventing or treating hyperproliferative vascular diseases in a mammal which comprises an amount of a compound of the formula I or II effective to inhibit intimal thickening in said mammal and a pharmaceutically acceptable carrier.
11. The composition of claim 10, which additionally contains an effective amount of a compound selected from the group consisting of rapamycin, mycophenolic acid, cyclosporin A, the substance FK 506 and brequimar sodium.



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention EP 95 10 0194 shall be considered, for the purposes of subsequent proceedings, as the European search report

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
P,X	TRANSPLANTATION PROCEEDINGS, vol. 26, no. 6, December 1994 pages 3244-3245, A.S. MACDONALD ET AL. 'EFFECTS OF LEFLUNOMIDE AND CYCLOSPORINE ON AORTIC ALLOGRAFT CHRONIC REJECTION IN THE RAT' * the whole document *	1-11	A61K31/42 A61K31/275
P,X	THE FASEB JOURNAL, vol. 8, no. 4, 15 March 1994 page A486 W. CAO ET AL. 'LEFLUNOMIDE, A NEW IMMUNOSUPPRESSANT, INHIBITS TYROSINE KINASE, CALCIUM SIGNALING AND DNA SYNTHESIS IN VASCULAR SMOOTH MUSCLE CELLS' * the whole document *	1-10	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
			A61K
INCOMPLETE SEARCH			
<p>The Search Division considers that the present European patent application does not comply with the provisions of the European Patent Convention to such an extent that it is not possible to carry out a meaningful search into the state of the art on the basis of some of the claims</p> <p>Claims searched completely : Claims searched incompletely : Claims not searched : Reason for the limitation of the search:</p> <p>see sheet C</p>			
Place of search		Date of completion of the search	Examiner
THE HAGUE		4 May 1995	Hoff, P
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number

EP 95 10 0194

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
P,X	THE JOURNAL OF HEART AND LUNG TRANSPLANTATION, vol. 13,no. 1, February 1994 page S68 C.R. GREGORY ET AL. 'LEFLUNOMIDE IS A NEW IMMUNOSUPPRESSANT THAT EFFECTIVELY REDUCES ARTERIAL INTIMAL THICKENING PRODUCED BY ALLOIMMUNE INJURY' * the whole document * ---	1-10	
P,X	EP-A-0 607 775 (HOECHST AKTIENGESELLSCHAFT) 27 July 1994 * abstract * * page 2, line 1 - page 3, line 24 * * claims * ---	1-10	TECHNICAL FIELDS SEARCHED (Int.Cl.6)
Y	WO-A-91 16892 (RORE INTERNATIONAL) 14 November 1991 * abstract * * page 1, line 1 - page 2, line 30 * * page 21, line 14 - line 26; claims; examples * ---	1-9	
Y	AMERICAN JOURNAL OF PHYSIOLOGY, vol. 260,no. 4, 1991 pages C721-C730, G.E. BILDER ET AL. 'TYRPHOSTINS INHIBIT PDGF-INDUCED DNA SYNTHESIS AND ASSOCIATED EARLY EVENTS IN SMOOTH MUSCLE CELLS' * the whole document * ---	1-9	
Y	FEBS LETTERS, vol. 334,no. 2, November 1993 pages 161-164, T. MATTAR ET AL. 'INHIBITION OF THE EPIDERMAL GROWTH FACTOR RECEPTOR TYROSINE KINASE ACTIVITY BY LEFLUNOMIDE' * the whole document * ---	1-9	
	---	-/--	



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number
EP 95 10 0194

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
Y	AGENTS AND ACTIONS, vol. 32,no. 1/2, 1991 pages 10-21, R.R. BARTLETT ET AL. 'LEFLUNOMIDE (HWA 486), A NOVEL IMMUNOMODULATING COMPOUND FOR THE TREATMENT OF AUTOIMMUNE DISORDERS AND REACTIONS LEADING TO TRANSPLANTATION REJECTION' * abstract * * page 17 - page 20 * ---	1-9	
X	TRANSPLANTATION PROCEEDINGS, vol. 25,no. 1, 1993 pages 745-746, J.W. WILLIAMS ET AL. 'IMMUNOSUPPRESSIVE EFFECTS OF LEFLUNOMIDE IN A CARDIAC ALLOGRAFT MODEL' * the whole document * ---	10,11	
D,X	EP-A-0 217 206 (HOECHST AKTIENGESELLSCHAFT) 8 April 1987 * the whole document * ---	10	TECHNICAL FIELDS SEARCHED (Int.Cl.6)
A	CURRENT OPINION IN CARDIOLOGY, vol. 8,no. 6, 1993 pages 938-950, M.L. FOEGH ET AL. 'MOLECULAR BIOLOGY OF INTIMAL PROLIFERATION' * abstract * * page 941 * * page 943 * * page 945 - page 946 * ---	1-9	
A	EP-A-0 551 182 (AMERICAN HOME PRODUCTS) 14 July 1993 * the whole document * -----	11	



EP 95 10 0194

- C -

Remark : Although claims 1-9 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.